

What to Consider in Your Practice

Who should be evaluated?

Any child or adolescent **age 4 to 18 years** who presents with 1) **academic or behavioral problems** and 2) symptoms of **inattention, hyperactivity, or impulsivity**. Children under 4 or over 12 years who first present with symptoms may have *Complex ADHD* and require special consideration. ADHD runs in families, therefore, particular early and frequent attention to children with a family history of ADHD is warranted. Genetics is a big *part* of the story, but not the whole story. For example, among identical twins, when one twin has ADHD, then there is a 55-90% chance that the other twin does too. The child of a parent with ADHD has a 25% chance of having ADHD as well.

Criteria and Diagnosis

- Diagnosis is based on [DSM-5 criteria](#) requiring a persistent pattern of inattention and/or hyperactivity–impulsivity that interferes with functioning or development.
- There are 4 dimensions/subtypes of ADHD identified in the DSM-5:

ADHD/I: primarily inattentive presentation	ADHD/HI: primarily hyperactive-impulsive presentation
ADHD/C: combined (inattentive + hyperactive-impulsive) presentation	ADHD other specified and unspecified ADHD

- Clinical Presentation—frequent patterns:
 - **Preschool age:** hyperactive; impulsive; inflexible; may be aggressive with peers
 - **Elementary school age:** struggles with listening in class; poor organizational skills; struggles with social interaction; difficulty functioning independently
 - **Adolescence:** struggles with learning, attention, and executive functioning; may find academic demands overwhelming
- The symptoms must occur in at least 2 settings (e.g., home, school, daycare). Information can be obtained from reports from parents/guardians, the child (if developmentally appropriate), teachers, other school personnel, and mental health clinicians who are involved in the child’s or adolescent’s care.
- A descriptive summary list of ADHD rating scales can be found at [CHADD’s Clinical Practice Tools](#). The NICHQ Vanderbilt Assessment Scale can be downloaded from the [AAP ADHD Resource Toolkit](#).

Sleep and ADHD

- Poor sleep quality or an inadequate amount of sleep can manifest as inattention, impulsiveness, hyperactivity, and oppositional behaviors in children without ADHD.
- Sleep problems also commonly occur in children with ADHD. One study revealed that children with ADHD have higher rates of daytime sleepiness compared to children without ADHD. Another study found that 50% of children with ADHD had signs of sleep-disordered breathing, compared to 22% of children without ADHD.
- It’s essential to evaluate for sleep behaviors when evaluating for ADHD and when providing ADHD management.

Treatment

Children ages 4-6 years

The **first line** of ADHD treatment is parent training in behavior management. If available, school-based behavioral interventions may also be implemented. When behavioral interventions do not provide significant improvement and the child continues to have serious problems, methylphenidate may be used.

Children and adolescents ages 6-18 years

The primary treatments include FDA-approved medications for ADHD along with parent training in behavior management and/or behavioral classroom interventions.

School-based behavioral interventions are typically designed and monitored through an Individualized Education Program (IEP) and/or a Section 504 plan.`

SPECIAL NEEDS INFORMATION AND RESOURCES

Local:

- Clark County Parent to Parent – 360-823-2347 x109
- Please join us in thanking and celebrating the retirement of our Clark County Children with Special Health Care Needs Nurse, Connie Callahan. Connie has played an important role in our community by providing care coordination services to families in our community. While we are sad to have Connie leave us, we are also happy to announce that we will be hiring a new nurse for this position.

Regional:	Advancing Integrated Mental Health Solutions (AIMS) Center	https://aims.uw.edu/targeted-condition/adhd
	WithinReach Family Health Hotline	1-800-322-2588, 1-800-833-6388 TTD http://www.parenthelp123.org/ Spanish: http://www.parenthelp123.org/es/
	Parent to Parent Support Programs of Washington	1-800-821-5927 https://arcwa.org/index.php/getsupport/parent_to_parent_p2p_programs
National:	Understood.org - Understanding ADHD in Your Child	https://www.understood.org/en/learning-thinking-differences/child-learning-disabilities/add-adhd/adhd-in-children
	AAP ADHD Resource Toolkit for Clinicians	https://www.aap.org/en-us/pubserv/adhd2/Pages/kit/data/assessdxframe.html
	U.S. Department of Education: Teaching Children With Attention Deficit Hyperactivity Disorder: Instructional Strategies and Practices	https://www2.ed.gov/rschstat/research/pubs/adhd/adhd-teaching_pg4.html
	Centers for Disease Control and Prevention (CDC) ADHD	https://www.cdc.gov/ncbddd/adhd/index.html
	Children and Adults with Attention-Deficit/Hyperactivity Disorder (CHADD)	https://chadd.org/ (general website) https://chadd.org/for-professionals/clinical-practice-tools/ (clinical practice tools)

References

- Alvarado, C., & Modesto-Lowe, V. (2017). Improving Treatment in Minority Children With Attention Deficit/Hyperactivity Disorder. *Clinical Pediatrics*, 56(2), 171–176. <https://doi.org/10.1177/000922816645517>
- Barbaresi WJ, Campbell L, Diekroger EA, Froehlich TE, Liu YH, O'Malley E, Pelham WE Jr, Power TJ, Zinner SH, Chan E. Society for Developmental and Behavioral Pediatrics Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents with Complex Attention-Deficit/Hyperactivity Disorder. *Journal of Developmental & Behavioral Pediatrics*. 2020;41:S35-S57. <https://doi.org/10.1097/DBP.0000000000000770>
- Centers for Disease Control (CDC). Data and Statistics About ADHD. <https://www.cdc.gov/ncbddd/adhd/data.html>
- Children and Adults with Attention-Deficit/Hyperactivity Disorder (CHADD). ADHD and Sleep Disorders. <https://chadd.org/about-adhd/adhd-and-sleep-disorders/>
- Coker, T. R., Elliott, M. N., Toomey, S. L., Schwebel, D. C., Cuccaro, P., Emery, S. T., Davies, S. L., Visser, S. N., & Schuster, M. A. (2016). Racial and Ethnic Disparities in ADHD Diagnosis and Treatment. *Pediatrics*, 138(3), 1–9. <https://doi.org/10.1542/peds.2016-0407>
- Lecturio Medical. (2018, March 14). Attention-Deficit/Hyperactivity Disorder (ADHD) – Pediatrics [Video]. <https://youtu.be/9ZUaDIxHqT4>
- Morgan, P. L., Staff, J., Hillemeier, M. M., Farkas, G., & Maczuga, S. (2013). Racial and Ethnic Disparities in ADHD Diagnosis From Kindergarten to Eighth Grade. *Pediatrics*, 132(1), 85–93. <https://doi.org/10.1542/peds.2012-2390>
- National Sleep Foundation. ADHD and Sleep. <https://www.sleepfoundation.org/articles/adhd-and-sleep>
- Wolraich ML, Hagan JF, Allan C, et al; Subcommittee on Children and Adolescents with Attention-Deficit/Hyperactive Disorder. Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents. *Pediatrics*. 2019;144(4):e20192528. (2020). *Pediatrics*, 145(3). <https://doi.org/10.1542/peds.2019-3997>